

Landscaping compromises for land degradation neutrality: The case of soil erosion in a Mediterranean agricultural landscape

Anna Smetanova, Stéphane Follain, Mélodie David, Rossano Ciampalini, Damien Raclot, Armand Crabit, Yves Le Bissonnais

▶ **To cite this version:**

Anna Smetanova, Stéphane Follain, Mélodie David, Rossano Ciampalini, Damien Raclot, et al.. Landscaping compromises for land degradation neutrality: The case of soil erosion in a Mediterranean agricultural landscape. Journal of Environmental Management, 2019, 235, pp.282-292. 10.1016/j.jenvman.2019.01.063 . hal-02067100

HAL Id: hal-02067100

https://institut-agro-dijon.hal.science/hal-02067100

Submitted on 21 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0301479719300696
Manuscript_6e5855dd4c90e41ced12b0f2596eba3e

Short Title: Target for catchment land degradation neutral management in Mediterranean

Long title: Landscaping compromises for land degradation neutrality: the case of soil erosion in a Mediterranean agricultural landscape

Anna Smetanová[a,b], Stéphane Follain[c], Mélodie David[b,d], Rossano Ciampalini[b], Damien Raclot[b,e], Armand Crabit[b], Yves Le Bissonnais[b]

[a] Research Group Ecohydrology and Landscape Evaluation, Ernst-Reuter PLatz 1, DE-10623, Berlin, Germany

[b] LISAH, INRA, IRD, Montpellier SupAgro, Univ Montpellier, FR-34060, Montpellier, France

[c] Agroécologie, AgroSup Dijon, INRA, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

[d] GEODE UMR 5602, Maison de la Recherche, FR-31500, Toulouse, France

[e] Institut Agronomique et Vétérinaire Hassan II, Dept. of Natural Resources and Environment, PO. Box. 6608, Madinat Al Irfane, Rabat, Morocco

corresponding author:

Anna Smetanová, Department of Ecohydrology and Landscape Evaluation, TU Berlin, Ernst-Reuter Platz 1, DE-10623, Berlin, Germany, anna.smetanova@gmail.com, Tel: +49 3031473534

© 2019 published by Elsevier. This manuscript is made available under the CC BY NC user license
https://creativecommons.org/licenses/by-nc/4.0/

Abstract

Soil erosion is the primary process driving land degradation. Using multiple scales of management to minimize soil erosion is crucial to achieve land degradation neutrality targets within the Sustainable Development Goals agenda. Land management (LM) influences both on-site and off-site erosion on the event-scale and over the long-term. However, each LM differs in effectiveness depending on the temporal scale considered. In order to understand how LM effects internal and external catchment dynamics, we apply LandSoil, a physically based landscape evolution model, to evaluate 7 LM scenarios over long- (30 years) and short-terms (event scale). LM scenarios included changes in land use and/or landscape structure. Under current LM, mean surface soil erosion was ~ 0.69 ± $39 \cdot 10^{-3}$ m over 30 years. In contrast, a single extreme event (435 mm/24h) in January resulted in ~ 0.62 ± $3 \cdot 10^{-3}$ m loss and ~ 0.04 ± $2 \cdot 10^{-3}$ m if it occurred in October. Heterogeneous patterns of erosion and deposition developed after 30 years, whereas extreme events dominantly showed soil loss and high catchment connectivity. Effectiveness of LM in erosion mitigation and sediment trapping differed according to temporal and spatial scales for each scenario. We concluded that multiple temporal and spatial scales must be incorporated in order to adaptively manage land degradation and meet neutrality targets.

Key words:

Land degradation, Degradation neutrality, Soil erosion, Landscaping, Mitigation strategy

1. Introduction

Land degradation neutrality targets of the Sustainable Development Goals (SDG 15.3) contain time-specific measures to avoid, reduce and reverse land degradation at both national and subnational levels (The Global Mechanism of UNCCD, 2016). Soil erosion by water is a major land degradation process (Orr et al, 2017). Therefore, land management seeking to minimize soil erosion is crucial for maintaining food and water security, climate regulation, soil ecosystem services (Dominati et al., 2010; Koch et al., 2013; Lal, 2004; McBratney et al., 2014;

Montanarella et al., 2016; and achieving of multiple SDGs (Keesstra et al., 2016; Orr et al., 2017).

It is expected, that soil erosion and suspended sediment yield in rivers will increase beginning in the mid-21st century due to climate change (Raclot et al., 2018; Routschek et al., 2014a, 2014b, Bussi et al., 2016; Rodríguez-Blanco et al., 2016). The majority of soil loss occurs during high-magnitude events of low frequency/recurrence, especially in Mediterranean environments where 3-10 of the largest daily events can account for over 50% of total soil loss (González-Hidalogo et al., 2007, 2010). Due to climate change, the frequency of high-magnitude rainfall events increases (e.g., Serpa et al., 2015). Therefore, considering seasonal patterns of soil loss (Smetanová et al., 2018) alongside water and sediment connectivity, and evaluating economic effects of on- and off-site effects of land management are all essential for preventing land degradation (García-Ruiz et al, 2017; Raclot et al., 2018).

Land use influences both soil erosion (sediment production) and sediment connectivity, while landscape structure can influence sediment transport and connectivity (Ciampalini et al., 2012; Coulthard & Van De Wiel, 2017; David et al, 2014; Follain et al., 2006; Fryirs, 2013). Modelling studies shown, that proper management can reduce the future effects of climate change on mid- and long-term soil erosion rates (Parroissien et al.2015; Rodriguez-Lloveras et al., 2016; Routschek et al., 2014a, 2014b). Modelling approaches successfully tested the effects of crop allocation, sediment trapping, and ditch optimisation techniques under mean or extreme rainfall events (David et al, 2014; Furlan et al., 2012; Gumiere et al., 2014; Levavasseur et al, 2016; Mullan et al., 2016; Nunes et al., 2013; Ronfort et al., 2011).

However, in order to meet land degradation neutrality targets, it is crucial to focus land management simultaneously on the relevant spatial and temporal scales for erosional processes (Larson et al., 1997; Stroosnijder, 2005) and stakeholders' scales of practice (Smetanová, et al, 2018). This requires providing of multi-foci, multi-scale solution. For example, a management reducing sheet and gully erosion during high-magnitude events of low

frequency/recurrence, and in long-term. Simultaneously, an effective management should reflect differing priorities of stakeholders operating in the same region. For example, the spatial scale of interests and mind-sets of a vineyards farmer and watershed manager might likely differ between understanding internal catchment dynamics on individual field scales (farmer) versus external catchment dynamics (e.g., sediment export to reservoirs). Furthermore, farmers establishing vineyards are interested in soil productivity over timescales that plants achieve highest productivity (10-30 years), while watershed managers might focus on reducing or preventing flash floods (single events). Understanding how land management effects on soil erosion contrast on temporal and spatial scales can help define targets for land management decisions aimed at reducing land degradation.

The objective of this study is to compare the internal and external catchment soil erosion dynamics for different land management scenarios over short-term (event scale) and long-term (30 years) scales, in order to answer the question: “Should management strategies for land degradation be tailored for specific spatial and temporal scales to achieve land degradation neutral management?”

2. Materials and Methods

2.1 Study site

The Roujan catchment (0.91 km^2), located in south France (43°30′N, 3°19′E) has been monitored for rainfall in the three meteorological stations equipped with tipping bucket rain gauges, and for continuous runoff and sediment concentration at parcel, mid- and whole catchment since 1992. It was described comprehensively by David et al. (2014). It belongs to a sub-humid Mediterranean climate (Peel et al., 2017) with mean annual rainfall 634 mm and a dry summer period, with maximum runoff and sediment yield from October to February. Elevation ranges 50m and slopes are 2–20% (David et al., 2014). Miocene marine and lacustrine sedimentary rocks with calcaric Regosols and Calcisols cover most of the catchment. Land use is dominated by vineyards (62%) followed by cereal and alfalfa production, and

scrubland (Guix-Hébrard et al., 2007). Chemical weeding in vineyards is performed using herbicide application alongside mechanical weeding by shallow (0.05–0.10 m) tillage with a duck foot cultivator in spring and autumn (Biarnès and Colin, 2006; Coulouma et al., 2006). Wheat and barley are rotated in the fields with annual crops. Fields are treated biannually in the post-harvest period (summer–autumn) with a disc-and-chisel plough down to depths of 7 to 12 cm. Landscape structure is characterised by 140 fields (0.6 ± 0.5 ha) with grass, bare soil or vegetated compacted soil strips along field borders, and an 11-km network of ditches (David et al. 2014). Current land use and landscape structures represent baseline land use scenario (Figure 1).

2.3. LandSoil model

LandSoil model was previously developed and calibrated in Roujan catchment (Ciampalini et al. 2012), and since then applied in variety of environments (e.g., Ciampalini et al.2011; Chartin et al, 2011; Lacoste et al., 2016). LandSoil is a spatial raster-based model for simulating water and tillage soil erosion as well as evolution of topography at plot to catchment scale. After modelling each event, LandSoil recalculates the elevation raster. Soil surface properties control water infiltration, runoff, and sediment concentration for each grid-cell and rainfall event (David et al., 2014; Leonard and Andrieux, 1998). Runoff in each grid cell is combination of input from upstream/upslope cells and runoff generated within the cell. Runoff flows in the flow line direction, or, if flow lines are interrupted, along liner landscape elements (roads, ditches, field boundaries, and tillage rows). The model uses a modified single-flow runoff model (Jenson and Domingue, 1988; Souchère et al., 1998) under eight possible flow directions. Both contour and downslope tillage were modelled in different fields with tillage transport coefficients spanning 111–139 kg m^{-1} (David et al., 2014).

2.4. Numerical experiment scheme

We considered two temporal dynamics (Figure 2):

(i) Long-term - 30 years. This period refers to long-term erosion rates over the timescale

that a new vineyard will reach peak productivity.

(ii) Single extreme rainfall event – maximum measured daily rainfall event. Events were modelled for January and October in order to consider variability in soil surface properties (David et al., 2014) that arise during periods when large erosion events usually occur (Smetanová et al., 2018).

We considered two spatial dynamics:

(i) Internal catchment dynamics – represented by fine spatial resolution modelling (1-meter grid) to simulate erosion and deposition patterns over the entire catchment. It related to spatial scale of farmers.

(ii) External catchment dynamics – represented by sediment export at the catchment outlet. It related to sediment input to river system, and management scale of watershed managers and policy makers.

We applied seven predictive land use scenarios that combine narrative and modelling methods developed by David et al. (2014). Three narratives were (i) stationary production (B - “baseline”, corresponding to current land mangement), (ii) more intensive production (I - “intensified”), and (iii) less intensive production (E - “extensified”) than stationary production. These narratives were transformed into seven scenarios (BLUS, ILU; ILS; ILUS; ELU, ELS, ELUS) by modifying the baseline land use and / or landscape structure by allocation rules for of land use (LU) and landscape structure (LS) as described in Figure 3.

2.5. Model inputs

Elevation and soil surface topography

LIDAR-based digital elevation model derived at 2-m resolution (David et al., 2014) represented initial input topography for all scenarios. Elevation was recalculated after every/each event by subtracting the depth of eroded soil or adding the height of deposited sediment in each raster cell. Soil erosion rates and sediment export (in $Mg \cdot km^{-2}$, where 1 $Mg \cdot km^{-2}$ corresponds to 0.01 tonnes per hectare) at the catchment outlet were recalculated at the end of each simulation period based on equations provided by David et al. (2014).

146

147 ***Rainfall dataset***

148 Long-term rainfall series were based on 10-years extension of empirical rainfall event
149 measurements in the Roujan catchment between 1992-2012 (David et al., 2014; Figure 4). The
150 total rainfall depth (mm), maximum intensity over 6 min (mm h–1) and rainfall duration (h) were
151 considered for each event, separated by at least a 6-hour dry period.
152 An event with return period >100 years was represented by extreme event measured in
153 Perpignan (120km from Roujan) on 26 October, 1915 (Cosadney and Robison, 2000; Meteo
154 France, 2018). Rainfall depth was 435 mm over 24 hours, but no measurement of rainfall
155 intensity was available. Rainfall intensity was estimated using the Montana law, with results
156 suggest that the Montana coefficient value exceeded the maximum intensity class for the
157 LandSoil model. Therefore, the maximum rainfall intensity class (>40 mm h^{-1}) was applied
158 based on calibration from Ciampalini et al. (2012). Model parameters and soil conditions are
159 described in additional detail by David et al. (2014).
160

161 ***Tillage Dataset***

162 Modelled tillage events occurred in fields with annual crops and in vineyards with mechanical
163 weeding. We simulated tillage twice a year in April or May and October. The exact day of a
164 tillage simulation was determined by cumulative rainfall depth (40 mm) since the last tillage
165 event.
166

167 3. Results

168 3.1 Temporal dynamics: Long-term vs. extreme event soil erosion
169

170 Under current land use and landscape structure conditions (BLUS), mean soil loss was 0.69 ±
171 $38.97 \cdot 10^{-3}$ m across the catchment over the 30-year simulation period (Table 1). Mean soil loss
172 here is the mean of all raster-cell values after 30 years of simulation. Each raster-cell value
173 represented cumulative elevation change after 30 years of simulation. Soil export at the outlet
174 (i.e., external catchment dynamics) under BLUS scenario was $9.02 \cdot 10^{-2}$ Mg·km^{-2}. Soil export at

175 catchment outlet varied from 4.74 (ELUS) to 119.20·10^{-2} Mg·km^{-2} (ILUS).
176 Erosional responses to single extreme rainfall events differed from long-term (30 years)
177 cumulative soil loss and were different for January and October. Mean soil loss in single
178 extreme rainfall event was 0.60 ± 2.72·10^{-3} m in January, and 0.00 ± 1.55·10^{-3} m in October
179 under BLUS scenario. Sediment export (external dynamics) was 16.4x higher in January than in
180 October. Similar patterns were observed in other scenarios, where mean soil loss for extreme
181 events were 1.3x (ILU) to 20x (ELS) higher in January relative to October.
182 Comparing mean soil loss of long-term and extreme events (Table 1) under BLUS scenario
183 showed that 8.3±14-times more soil was eroded after thirty years than by a single extreme
184 event in January.
185 Sediment exported by a single event in January or October is only reached as the cumulative
186 effect of many erosion events under normal rainfall conditions. This comparison could be called
187 'equal erosion delivery', which refers to the duration required for continuous "normal" erosion to
188 match the quantity of sediment delivered by an extreme event. In BLUS, sediment exported in
189 January was reached after ~27 years, while sediment exported during October was matched
190 after ~2 years. Sediment exported in all January extreme events was reached in ~21-26 years
191 under extensified land use scenarios, but dropped drastically to~2-4 years if October extreme
192 events were considered. Under ILU and ILUS, soil loss by both extreme events were more
193 similar than by all remaining scenarios. Under long-term rainfall conditions, an equivalent
194 amount of sediment export was reached in 6-7 years.
195
196 3.2 Internal catchment dynamics
197
198 Long-term rainfall conditions under current land use and landscape structure (BLUS) led to soil
199 redistribution within the catchment (Figure 5B). Soil loss ranged from 0.01 to 0.3 m in vineyards
200 with chemical weeding, with less than 0.01 m loss in scrubland and no tillage annual crop fields.
201 In vineyards with mechanical weeding and in fields with tilled annual crops, erosion ranged from
202 0.01 to 0.3 m and deposition >0.01 m occurred. Rills created in parcels with no tillage and
203 scrubland were infilled by shallow deposition (<0.01 m). Zero net erosion or deposition (<10^{-12}m)

204 was observed on or along linear landscape structures (e.g., roads, field borders).
205
206 In contrast to previous results, during an extreme event in January (Figure 5C), net soil loss was
207 0.01 to 0.3 m in all land uses excluding scrubland, where a series of long rills with depths <
208 0.01 m, were created. Rills deeper than 0.3 m were formed along some recent and/or historical
209 linear landscape structures. Deposition > 0.1 m only took place on some of the vegetated strips.
210 During an extreme event in October, mean soil loss was lower than that from extreme events in
211 January (Table 1). Furthermore, the rill network within scrubland and along field borders was
212 less dense and generally, shorter rills infilled with less than 0.01 m sediment (Figure 5D). “Equal
213 erosion delivery” duration was ~27-years for an extreme event in January. However, soil
214 redistribution patterns were completely different (Figure 5E). Sediment exported during an
215 extreme rainfall event in October was reached after ~2 years of long-term rainfall conditions
216 (Figure 5F), but contrary to extreme rainfall in October (Figure 5D) nearly zero erosion or
217 deposition occurred within most of the parcels under normal rainfall conditions.
218
219 For other scenarios, the relation between long-term and extreme event internal catchment
220 dynamics was similar to those in BLUS (Supplementary Info 1). Normal rainfall simulations led
221 to heterogeneous soil redistribution patterns with areas of both erosion and deposition after 30
222 years, while nearly zero net soil erosion over the majority of the catchment during extreme
223 events.
224 Differences in internal catchment dynamics for each temporal scale are shown using an
225 example of upper (terraced) slopes divided to two parcels (Figure 6). The upper parcel was
226 divided to one (ILS) or more (BLUS, ELS) fields with mechanical weeding. Chemical weeding
227 was applied in ILU and ILUS scenarios, while scrubland covered the parcel in ELU and ELUS
228 (Figure 6A). The lower parcel was covered by scrubland in all scenarios, but the field border
229 cover varied (Figure 6A). Under scenarios of intensified land use (ILU and ILUS), 0.01-0.3 m of
230 soil was removed from surface in vineyards with mechanical weeding. Furthermore, deep rills
231 (0.01-0.3 m) in scrublands were infilled with transported material after 30 years (Figure 6B).
232 During extreme events in January (Figure 6C), the scrubland rill network was much less dense

compared to long-term simulations. However, during extreme event in January some of the partly infilled rills cut into adjacent field borders and continued eroding. Additionally, exclusively soil erosion occurred downhill of the scrubland field border during extreme events in January. Substituting vegetated pathways with bare soil (Figure 6A-ILS) increased connectivity between parcels and concentration of overland flow from upper parcel in lower parcels under long-term rainfall conditions (Figure 6B-ILS). During extreme events (Figure 6C, 6D –ILS) erosion <0.01 m occurred along some rills in scrubland. Under ILS, equal erosion delivery was attained after 17 years in comparison to January extreme event. Soil redistribution after 17 and 30-years under ILS was similar. For ILU and ILUS, equal erosion delivery was 7 and 6 years. Again the soil erosion pattern of equal soil erosion delivery was similar to long-term soil redistribution pattern (Figure 6E –ILU and ILUS).

Under ELU overland flow generated in upper parcel (scrubland, Figure 6A-ELU) caused less intensive erosion on-site than vineyards parcels in stationary and more intensive production scenarios in 30 years (Figure 6B). Furthermore, scrubland led to rill formation and their infilling in lower parcel after 30 years (Figure 6B-ELU). In ELUS, combined land use and landscape structure changes helped protect rill incision (Figure 6B ELUS), but as Figure 6B ELS shows, change of landscape structure alone was not sufficient to prevent rill erosion. Under extreme events, erosion was low along scrubland field borders, and downslope rills emerged only under ELS. During the year when sediment export from long-term rainfall conditions equalled extreme events, soil redistribution patterns was more related to long-term precipitation patterns (Figure 6, Supplementary Info 1).

3.3 External catchment dynamics

Sediment export is referred to external catchment dynamics, and is reported in section 3.1, Table 1. In scenarios with intensified land use (ILU, ILUS), total sediment export over long-term conditions was over 10-fold higher than BLUS, over 4-fold higher than ILS. Total sediment export under extreme event in January was in ILU and ILUS over 2.6-fold higher than BLUS, and over 1.6-fold higher than ILS. For scenarios with less intensive production than BLUS, long-term sediment export at catchment outlet by 10% (ELU), 40% (ELS) and 50% (ELUS) in

comparison with BLUS sediment export.

Sediment export under all scenarios was increasingly stable over the first four years of model simulations, followed by sudden increases in the fifth year triggered by rainfall (Figure 7). The sediment export response to natural rainfall conditions differed between the modelled scenarios. Intensified scenarios responded with higher sediment production (i.e., erosion) and transport. Soil export in ILUS more than doubled, and increased ~1.5-fold in ILU and ILS (Figure 7B). On the contrary, ELU exhibited increased sediment export than ELS. Threshold behaviours with different magnitudes of sediment response existed for all scenarios in several years (e.g., year 16, year 24).

Figure 7B compares sediment export by extreme events in January with long-term sediment export. Intensified scenarios reached the sediment export of BLUS within first 4 to 10 years (for ILUS and ILS, respectively).

4. Discussion

4.1 Land management optimisation: understanding the impact on differing temporal scales

Comparing scenarios over different time scales showed that both external and internal catchment dynamics differed for long-term and extreme-event rainfall conditions. Mean and variability of soil loss were greater after long-term rainfall events than extreme rainfall conditions. However, sediment export at the catchment outlet from a single extreme event (in January) accounted for 89% (BLUS) of long-term sediment export. Within-catchment and within-parcels erosional hotspots formed over the 30 year simulations. Hotspots' spatial distribution differed for each land use scenario, while landscape structure influenced within-catchment (dis-) connectivity. Soil redistribution within parcels was strongly influenced by erosion from tillage (David et al, 2014), which temporarily influences soil properties and micro-landforms (e.g., infilling rills, creating tillage rows) and thus, hydrology and associated sediment connectivity along hillslopes.

291 During extreme events, most of the catchment was connected. Soil erosion removed up to 0.01-
292 m from almost the entire catchment independent of land use or landscape structure (except for
293 scrubland and some of the filed borders, grass strips, pathways). Land uses or landscape
294 structures sometimes acted as barriers or buffers under normal rainfall conditions, but were less
295 effective in damping overland flow and sediment trapping. Yet, effects of single extreme rainfall
296 events differed for October and January depending on soil-vegetation properties in the
297 respective month. In October, transport-limited deposition took place in some pathways,
298 whereas only transport and zero net erosion or deposition took place in January. Antecedent
299 conditions such as previous rainfall or tillage influenced sensitivity of sediment response to
300 seasonal changes in soil hydrological properties, demonstrated previously by monitoring studies
301 (e.g., Biddocu et al., 2017; Inoubli et al., 2017; Raclot et al., 2009, Smetanova et al., under
302 revision).
303
304 4.2 Land management optimisation: understanding effects of management measures under
305 different scenarios
306
307 We discuss the effect of land management on land-degradation based on area-specific
308 sediment budgets for both long-term and extreme events (Figure 8).
309 Chemical weeding led to erosion when applied on vineyards in all scenarios and timescales.
310 Mechanical weeding, was also a net source of sediment under extreme rainfall conditions in
311 January, but over 30 years, had nearly neutral sediment budget. According to David et al.
312 (2014), mechanical weeding caused three-times less sediment erosion compared to chemical
313 weeding over 100 years. However, extensive mechanical weeding in ELU without using
314 extensive landscape structures (such as in ELS) increased sediment export more than both
315 current land use and extensive landscape structure in ELS (Table 1). This confirms that land
316 use and landscape structure management must be used in combination to effectively decrease
317 both sediment production (i.e., erosion) from farmers' fields and sediment exported from the
318 catchment to river system.
319 Grass strips and vegetated strips were the most effective management strategies for trapping

sediment at both temporal scales (and in 100 years as in David et al., 2014), but the sediment trapped was highest for extreme events. Total area and spatial distribution of vegetated strips was identical under BLUS; ILU and ILUS, but sediment trapping in vegetated strips decreased with decreasing sediment production. Sediment trapping efficiency of vegetated strips differed with (i) amount of sediment produced, (ii) spatial redistribution of landscape elements and land uses, and (iii) single (process) versus cumulative events (long-term processes). Trapping efficiency was directly linked to sediment connectivity during extreme events and over the long-term. This confirm previous findings that both sediment connectivity and sediment trapping efficiency are dependent on spatial distribution of management (Collin et al., 2012; Gumiere et al., 2011; Mekonnen et al., 2015; Ramos et al., 2010). Using a raster-based approach allowed us to assess sediment production and transport at individual parcel scale at both temporal scales (example in Figure 6). However, we did not calculate soil loss and effectiveness of measures for each farmer at different time scales, which is possible and recommended for supporting decision-making schemes (Souchère et al., 2010). At the catchment scale, soil surface properties drove sediment export regardless of land use (David et al., 2014). Therefore, management techniques aimed at the soil surface can reduce soil erosion. Such approaches include straw mulch or intercropping, and are known to decrease soil erosion under long-term and extreme rainfall conditions (Biddocu et al., 2014, 2016, 2017; Blavet et al., 2009; Cerdà et al., 2016; Gómez et al., 2009, 2014; Kosmas et al., 1997; Prosdocimi et al., 2016; Raclot et al., 2009; Ramos et al., 2007; Rodrigo-Comino et al., 2016; Ruiz-Colmenro et al., 2011). Management strategies can be either (i) permanently incorporated in land use scenarios, or (ii) used in conjunction with others (e.g., geotextiles, mobile sediment trapping barriers) as ad-hoc measures during extreme events (Sherrif et al., 2016).

4.3 Pathways for applicable optimisation for land-degradation neutral management

Our results suggest that using multiple temporal, spatial and management scales to evaluate soil erosion and connectivity can significantly improve existing modelling approaches (Bussi et al., 2016; Collin et al., 2012; David et al., 2014; Gumiere et al., 2014; Mullan et al., 2016; Nunes

et al., 2013; Paroissien et al, 2015; Ramos et al., 2015; Rodriguez-Lloveras et al., 2016; Ronfort et al., 2011; Routschek et al., 2014). Our approach includes temporally- and spatially-adjusted measures that are applicable for meeting soil erosion and land degradation neutrality targets (The Global Mechanism of UNCCD, 2016). We demonstrate that proposing land degradation neutrality measures in agricultural catchments is extremely complex due to (i) natural variability of catchment responses to normal rainfall conditions and extreme events, and (ii) variability in cumulative response depending on the spatial distribution of selected measures. Furthermore, we showed that (iii) the same sediment export (sediment yield) values represented different patterns of soil erosion and deposition within a catchment. The complexity of applying measures to reduce land degradation is reinforced by differing interests and management aims of each stakeholder (e.g., farmer or watershed manager; Smetanová et al., 2018). Therefore, modelling approaches applied on differing spatial and temporal scales are valuable aid for participative decision making on best practice management (Hewett et al., 2018, Keesstra et al., 2018).

5. CONCLUSION

Our research proved, that tailoring specific management strategies for specific spatial and temporal scales might be suitable in order to achieve land degradation neutral management. Our research contributions included the following in regards to attaining land degradation neutrality goals:

1. Managing changes to land use or cover won't appropriately serve land degradation neutrality targets at the catchment scale (sub-national level). It is essential to include landscape structure change and connectivity management into land use scenarios.
2. Internal and external catchment dynamics differ for long-term and extreme events, and therefore multiple temporal, spatial, and management scales must be compared and incorporated in to planning adaptive management for land degradation. Optimally, this will span from farm- to catchment-scales, as well as sub-national levels.
3. Modelling approaches such as LandSoil enable comparison of management efficiency for both long- and extreme event-scales. Additionally, the model improves selection of

timely and adaptive management scenarios best serving land degradation neutrality targets (SDGs 15.3)

We suggest that holistic land degradation neutrality management can be selected by using multi-scale consideration based on models incorporated in participative decision making. Using such inputs will bring multiple co-benefits fostering life on land, including avoiding, reducing and reversing land degradation processes.

Acknowledgement

This work benefits from the financial support of ANR-08-VULN-006 (LANDSOIL project) and MESOEROS21 (ANR-06-VULN-012), ORE OMERE, MISTRALS/Sicmed R_Osmed and MASCC (funded by ARIMNET2). Anna Smetanová received Marie-Curie FP7 COFUND People Programme AgreenSkills fellowship (grant 267196). Author thank to Dr. Mitchell Donovan for his helpful comments and remarks.

References

Biarnès A, Colin F. 2006. Methodology for assessing the hydrological impact of weed control practices with a view to management of Mediterranean vine growing catchments. International Journal of Sustainable Development 9, 161–179. doi: 10.1504/IJSD.2006.011639

Biddocu M, Ferraris S, Pitacco A, Cavallo E. 2017. Temporal variability of soil management effects on soil hydrological properties, runoff and erosion at the field scale in a hillslope vineyard, North-West Italy. Soil & Tillage Research 165, 46–58.

Biddoccu M, Ferraris S, Opsi F, Cavallo E. 2016. Long-term monitoring of soil management effects on runoff and soil erosion in sloping vineyards in Alto Monferrato (North-West Italy). Soil & Tillage Research 155, 176–189. doi:10.1016/j.still.2015.07.005.

Biddoccu M, Opsi F, Cavallo E. 2014. Relationships between runoff and soil losses with rainfall characteristics and a long-term soil management practices in a hilly vineyard (Piedmont, NW Italy). Soil Science and Plant Nutrition 60, 92–99. doi: 10.1080/00380768.2013.862488.

Blavet D, De Noni G, Le Bissonnais Y, Leonard M, Maillo L, Laurent JY, Asseline J, Leprun JC,

Arshad MA, Roose E. 2009. Effect of land use and management on the early stages of soil water erosion in French Mediterranean vineyards. Soil & Tillage Research 106, 124–136.

Bussi G, Dadson SJ, Prudhomme C, Whitehead P. 2016. Modelling the future impacts of climate and land-use change on suspended sediment transport in the River Thames (UK). Journal of Hydrology 542, 357-372. doi: 10.1016/j.jhydrol.2016.09.010

Cerdà A, González-Pelayo O, Giménez-Morera A, Jordán A, Pereira P, Novara A, Brevik EC, Prosdocimi M, Mahmoodabadi M, Keesstra S, García Orenes F, Ritsema C. 2016. The use of barley straw residues to avoid high erosion and runoff rates on persimmon plantations in Eastern Spain under low frequency — high magnitude simulated rainfall events. Soil Research 54, 154-165.

Ciampalini R, Billi P, Follain S, Le Bissonnais Y, Ferrari G, Borselli L. 2011. Long-term soil erosion analysis on the terrace systems of the archaeological area of Aksum (Ethiopia) (1800 BC – today). IAG Regional meeting (Addis Abeba).

Ciampalini R, Follain S, Le Bissonnais Y. 2012. LandSoil: a model for the analysis of erosion impact on agricultural landscape evolution. Geomorphology 175, 25–37. doi: 10.1016/j.geomorph.2012.06.014

Ciampalini R, Follain S, Cheviron B, Le Bissonnais Y, Couturier A, Walter C. 2017. Local sensitivity analysis of the LandSoil erosion model applied on a virtual catchment". In: Srivastava P, Petropoulos G. (Eds.) Sensitivity Analysis in Earth Observation Modelling, Elsevier, pp 55-73.

Chartin C, Bourennane H, Salvador-Blanesa S, Hinschberger F, Macaire J-J. 2011. Classification and mapping of anthropogenic landforms on cultivated hillslopes using DEMs and soil thickness data — Example from the SW Parisian Basin, France. Geomorphology 135, 8-20. doi: 10.1016/j.geomorph.2011.07.020

Collin F, Moussa R, Louchart X. 2012. Impact of the spatial arrangement of land management practices on surface runoff for small catchments. Hydrological Processes 26, 255-271. doi: 10.1002/hyp.8199

Cosandey C, Robinson M. 2000. Hydrologie continentale. Armand Colin, Paris, 2000, 368 p.

Coulouma G, Boizard H, Trotoux G, Lagacherie P, Richard G. 2006. Effect of deep tillage for vineyard establishment on soil texture: a case study in southern France. Soil & Tillage Research

436 88, 132–143. doi: 10.1016/j.still.2005.05.002
437 Coulthard TJ, Van De Wiel MJ. 2017. Modelling long term basin scale sediment connectivity,
438 driven by spatiall and land use changes. Geomorphology 277, 265-281. doi:
439 10.1016/j.geomorph.2016.05.027
440 David M, Follain S, Ciampalini R, Le Bissonnais Y, Couturier A, Walter C. 2014. Simulation of
441 medium-term soil redistributions for different land use and landscape design scenarios within a
442 vineyard landscape in Mediterranean France. Geomorphology 214, 10-21. doi:
443 10.1016/j.geomorph.2014.03.016
444 Dominati E, Patterson M, Mackay A. 2010. A framework for classifying and quantifying the
445 natural capital and ecosystem services of soils. Ecological Economics 69, 1858–1868. doi:
446 10.1016/j.ecolecon.2010.05.002
447 Follain S, Minasny B, McBratney AB, Walter C. 2006. Simulation of soil thickness evolution in a
448 complex agricultural landscape at fine spatial and temporal scales. Geoderma 133, 71-86.
449 Furlan A, Poussin J-C, Mailhol J-C, Le Bissonnais Y, Gumiere SJ. 2012. Designing
450 management options to reduce surface runoff and sediment yield with farmers: An experiment
451 in south-western France. Journal of Environmental Management 96, 74-85.
452 Fryirs K. 2013. (Dis)Connectivity in catchment sediment cascades: a fresh look at the sediment
453 delivery problem. Earth Surface Processes & Landforms 38, 30–46. doi: 10.1002/esp.3242
454 García-Ruiz JM, Beguería S, Lana-Renault N, Nadal-Romero E, Cerda A. 2017. Ongoing and
455 emerging questions in water erosion studies. Land Degradation & Development 28, 5-21.
456 Gómez JA, Sobrinho TA, Giráldez JV, Fereres E. 2009. Soil management effects on runoff,
457 erosion and soil properties in an olive grove of Southern Spain. Soil & Tillage Research. 102, 5–
458 13. doi:10.1016/j.still.2008.05.005.
459 Gómez JA, Vanwallenghem T, De Hoces A, Taguas EV. 2014. Hydrological and erosive
460 response of a small catchment under olive cultivation in a vertic soil during a five-year period:
461 implications for sustainability. Agric. Ecosyst. Environ. 188, 229–244.
462 Guix-Hébrard N, Voltz M, Trambouze W, Garnier F, Gaudillère JP, Lagacherie P. 2007.
463 Influence of watertable depths on the variation of grapevine water status at the landscape scale.

European Journal of Agronomy 27, 187–196. doi: 10.1016/j.eja.2007.03.005

Gumiere SJ, Bissonnais Y, Raclot D, Cheviron B. 2011. Vegetated filter effects on sedimentological connectivity of agricultural catchments in erosion modelling: A review. Erath Surface Processes and Landforms 36, 3-19.

Gumiere SJ, Bailly J-S, Cheviron B, Raclot D, Le Bissonnais Y, Rousseau A. 2014. Evaluating the impact of the spatial distribution of land management practices on water erosion: case study of a Mediterranean catchment. Journal of Hydrology Engineering. doi: 10.1061/(ASCE)HE.1943-5584.0001076, C5014004.

Hewett C, Simpson C, Wainwright J, Hudson S. 2018. Communicating risk to infrastructure due to soil erosion: a bottom up approach. Land Degradation & Development 29, 1282–1294.

Inoubli N, Raclot D, Moussa R, Habaieb H, Le Bissonnais Y. 2016. Soil cracking effects on hydrological and erosive processes in Mediterranean cultivated vertisols. Hydrological Process, 30, 4154-4167.

Jenson SK, Domingue JO. 1988. Extracting topographic structure from digital elevation data for geographic information system analysis. Photogrammetric Engineering and Remote Sensing 54, 1593-1600.

Keesstra SD, Bouma J, Wallinga J, Tittonell P, Smith P, Cerdà A, Montanarella L, Quinton JN, Pachepsky Y, van der Putten WH, Bardgett JD, Moolenaar S, Mol G, Jansen B, Fresco, LO. 2016. The significance of soils and soil science towards realization of the United Nations Sustainable Development Goals. Soil 2, 111–128. doi: 10.5194/soil-2-111-2016

Keesstra SD, Nunes JP, Novara A, Finger D, Avelar D, Kalantari Z, Cerda A. 2018. The superior effect of nature based solutions in land management for enhancing ecosystem services. Science of the Total Environment 610-611, 997-1009. doi: 10.1016/j.scitotenv.2017.08.077

Koch A, Field D, McBratney A B, Adams M, Hill R, Craw-ford J, Minasny B, Lal R, Abbott L, O'Donnel A, Angers D, Baldock J, Barbier E, Binkley D, Parton W, Wall DH, Bird M, Chenu C, Flora CB, Goulding K, Grunwald S, Hempel J, Jastrow J, Lehmann J, Lorenz K, Morgan CL, Whitehead D, Young I, Zimmermann M. 2013. Soil security: solving the global soil crisis. Global Policy 4, 434–441. doi: 0.1111/1758-5899.12096

493 Kosmas C, Danalatos N, Cammeraat LH, Chabart M, Diamantopoulos J, Farand R, Gutierrez L,
494 Jacob A, Marques H, Martinez-Fernandez J, Mizara A, Moustakas N, Nicolau JM, Oliveros C,
495 Pinna G, Puddu R, Puigdefabregas J, Roxo M, Simao A, Stamou G, Tomasi N, Usai D, Vacca
496 A. 1997. The effect of land use on runoff and soil erosion rates under Mediterranean conditions.
497 CATENA, 29, 45–59.
498 Lacoste M, Viaud V, Michot D, Walter C. 2016. Model-based evaluation of impact of soil
499 redistribution on soil organic carbon stocks in a temperate hedgerow landscape. Earth Surface
500 Processes and Landforms, 41, 1536-1549. doi: 10.1002/esp.3925
501 Lal R. 2004. Soil Carbon Sequestration Impacts on Global Climate Change and Food Security.
502 Science 304, 1623–1627. doi: 10.1126/science.1097396
503 Leonard J, Andrieux P. 1998. Infiltration characteristics of soils in Mediterranean vineyards in
504 southern France. Catena 32, 209–223. doi: 10.1016/S0341-8162(98)00049-6
505 Levavasseur F, Bailly J-S, Lagacherie P. 2016. Are ditch networks optimised for mitigating rill
506 erosion in cultivated Mediterranean landscapes? A numerical experiment. Land Use Policy 50,
507 441-448.
508 Meteo France, 2018. Pluies extrêmes en France métropolitaine.
509 http://pluiesextremes.meteo.fr/france-metropole/Fantastique-chute-de-pluie-sur-Perpignan.html.
510 Accessed: 15 May 2018.
511 McBratney AB, Field DJ, Koch A. 2014. The dimensions of soil security. Geoderma 213, 203–
512 213. doi: 10.1016/j.geoderma.2013.08.013
513 Mekonnen M, Keesstra SD, Stroosnijder L, Baartman JEM, Maroulis J. 2015. Soil conservation
514 through sediment trapping: A review. Land Degradation & Development 26, 544–556.
515 Montanarella L, Pennock DJ, McKenzie N, Badraoui M, Chude V, Baptista I, Mamo T,
516 Yemefack M, Singh Aulakh M, Yagi K, Young Hong S, Vijarnsorn P, Zhang G-L, Arrouays D,
517 Black H, Krasilnikov P, Sobocká J, Alegre J, Roberto Henriquez C, de Lourdes Mendonça-
518 Santos M, Taboada M, Espinosa-Victoria D, Al Shankiti A, Alavi Panah SK, El Mustafa Elsheikh
519 EA, Hempel J, Camps Arbestain M, Nachtergaele F, Vargas R. 2016. World's soils are under
520 threat. Soil 2, 79-82. doi: 10.5194/soil-2-79-2016
521 Mullan D, Vandaele K, Boardman J, Meneely J, Crossley LH. 2016. Modelling the effectiveness

of grass buffer strips in managing muddy floods under a changing climate. Geomorphology 102–120. doi: 10.1016/j.geomorph.2016.07.012

Nunes JP, Seixas J, Keizer JJ. 2013. Modelling the response of within-storm runoff and erosion dynamics to climate change in two Mediterranean watersheds: A multi-model, multi-scale approach to scenario design and analysis. Catena 102, 27-39. doi: 10.1016/j.catena.2011.04.001

Orr BJ, Cowie AL, Castillo Sanchez VM, Chasek P, Crossman ND, Erlewein A, Louwagie G, Maron M, Metternicht GI, Minelli S, Tengberg AE, Walter S, Welton S. 2017. Scientific Conceptual Framework for Land Degradation Neutrality. A Report of the Science-Policy Interface. United Nations Convention to Combat Desertification (UNCCD), Bonn, Germany. ISBN 978-92-95110-42-7 (hard copy), 978-92-95110-41-0 (electronic copy).

Paroissien J-B, Darboux F, Devillers B, Mouillot F, Raclot D, Le Bissonnais Y. 2015. A method for modelling the effects of climate and land use changes on erosion and sustainability of soil in a Mediterranean watershed (Languedoc, France). Journal of Environmental Management 150, 57-68. doi: 10.1016/j.jenvman.2014.10.034

Peel, M.C., Finlayson, B.L., McMahon, T.A. 2007. Updated world map of the Köppen-Geiger climate classification. Hydrology & Earth System Sciences, 11, 1633–1644.

Prosdocimi, M., Cerdà, A., Tarolli, P., 2016. Soil water erosion on Mediterranean vineyards: a review. Catena 141 (2016), 1–21.

Raclot D, Le Bissonnais Y, Annabi M, Sabir M, Smetanova A. 2018. Main issues for preserving Mediterranean soil resources from water under global change. Land Degradation & Development 29, 789–799.

Raclot D, Le Bissonnais Y, Louchart Y, Andrieux P, Moussa R, Voltz M. 2009.Soil tillage and scale effects on erosion from fields to catchment in a Mediterranean vineyard area. Catena 66, 198-210.

Ramos MC, Martínez-Casasnovas JA. 2007. Soil loss and soil water content affected by land levelling in Penedès vineyards, NE Spain. Catena 71, 210–217.

Ramos MC, Martínez-Casasnovas JA. 2010. Effects of field reorganisation on the spatial variability of runoff and erosion rates in vineyards of Northeastern Spain. Land Degradation &

551 Development 21, 1–12.
552 Ramos MC, Benito C, Martínez-Casasnovas JA. 2015. Simulating soil conservation measures
553 to control soil and nutrient losses in a small, vineyard dominated, basin. Agriculture,
554 Ecosystems & Environment, 213, 194–208.
555 Rodrigo-Comino J, Iserloh T, Morvan X, Malam Issa O, Naisse C, Keesstra SD, Cerdà A,
556 Prosdocimi M, Arnáez J, Lasanta T, Concepión Ramos M, Marqués MJ, Ruiz-Colmenero M,
557 Bienes R, Ruiz Sinoga JD, Seeger M, Ries JB.2016. Soil Erosion Processes in European
558 Vineyards: A Qualitative Comparison of Rainfall Simulation Measurements in Germany, Spain
559 and France. Hydrology 3, 6. doi:10.3390/hydrology3010006
560 Rodríguez-Blanco ML, Arias R, Taboada-Castro MM, Nunes JP, Keizer JJ, Taboada-Castro T.
561 2016. Potential Impact of Climate Change on Suspended Sediment Yield in NW Spain: A Case
562 Study on the Corbeira Catchment. Water 8, 444. doi: 10.3390/w8100444
563 Rodriguez-Lloveras X, Buytaert W, Benito G. 2016. Land use can offset climate change induced
564 increases in erosion in Mediterranean watersheds. Catena 143, 244-255. doi:
565 /10.1016/j.catena.2016.04.012
566 Ronfort C, Souchère V, Martin P, Sebillotte C, Castellazzi MS, Barbottin A, Meynard JM, Laignel
567 B. 2011. Methodology for land use change scenario assessment for runoff impacts: A case
568 study in a north-western European Loess belt region (Pays de Caux, France). Catena 86, 36–
569 48. doi:10.1016/j.catena.2011.02.004
570 Routschek A, Schmidt J, Enke W, Deutschlander T. 2014a. Future soil erosion risk — Results of
571 GIS-based model simulations for a catchment in Saxony/Germany. Geomorphology 206, 299-
572 306. doi: 10.1016/j.geomorph.2013.09.033
573 Routschek A, Schmidt J, Kreienkamp F. 2014b. Impact of climate change on soil erosion — A
574 high-resolution projection on catchment scale until 2100 in Saxony/Germany. Catena 121, 99-
575 109. doi: 10.1016/j.catena.2014.04.019
576 Serpa D, Nunes JP, Santos J, Sampaio E, Jacinto R, Veiga S, Lima JC, Moreira M, Corte-Real
577 J, Keizer JJ, Abrantes N. 2015. Impacts of climate and land use changes on the hydrological
578 and erosion processes of two contrasting Mediterranean catchments. Science of Total
579 Environment 538, 64–77. doi: 10.1016/j.scitotenv.2015.08.033

580 Sherriff SC, Rowan JC, Fenton O, Jordan P, Melland AR, Mellander P-E, Ó hUallacháin D.
581 2016. Storm Event Suspended Sediment-Discharge Hysteresis and Controls in Agricultural
582 Watersheds: Implications for Watershed Scale Sediment Management. Environmental Sciences
583 & Technology, 50, 1769-1778. doi: 10.1021/acs.est.5b04573

584 Smetanová A, Paton E, Maynard C, Tindale S, Fernandez-Getino A-P, Marques MJ, Bracken L,
585 Le Bissonnais Y, Keesstra S. 2018. Stakeholders' perception of the relevance of water and
586 sediment connectivity in water and land management. Land Degradation & Development, doi:
587 10.1002/ldr.2934

588 Smetanová A, Le Bissonnais Y, Raclot D, Nunes JP, Licciardello F, Le Bouteiller C, Latron J,
589 Rodríguez-Caballero E, Mathys N, Klotz S, Mekki I, Gallart F, Solé Benet A, Pérez Gallego N,
590 Andrieux P, Moussa R, Planchon O, Marisa Santos J, Alshihabi O, Chikhaoui M., Follain S.,
591 2018. Patterns of temporal variability and time compression of sediment yield in small
592 Mediterranean catchments. Soil Use & Management 34, 388-403. doi: 10.1111/sum.12437.

593 Souchère V, King D, Daroussin J, Papy F, Capillon A. 1998. Effects of tillage on runoff
594 directions: consequences on runoff contributing area within agricultural catchments. Journal of
595 Hydrology 206, 256-267.

596 Souchère V, Millair L, Echeverria J, Bousquet F, Le Page C, Etienne M. 2010. Co-constructing
597 with stakeholders a role-playing game to initiate collective management of erosive runoff risks at
598 the watershed scale. Environmental Modelling & Software 25, 1359–1370.

599 Stroosnijder, L. 2005. Measurement of erosion: Is it possible? CATENA, 64, 162-173

600 The Global Mechanism of UNCCD. 2016. Achieving Land Degradation Neutrality at the Country
601 Level: Building Blocks for LDN Target Setting. Bonn, Germany. ISBN 978-92-95110-31-1

Figure description:

Figure 1. Study area (A), land use (B), and landscape structures (C)

Figure 2: Flowchart of numerical experiment in LandSoil. Model inputs were digital elevation model, dynamic soil properties, seven land management scenarios, and rainfall. Two temporal (long-term, single event) and two spatial dynamics (internal and external) were considered. Red triangle indicates management priorities of a farmer, inverted red triangle priorities of a watershed manager. Outputs related to internal (v) and external (inverted v) catchment dynamics were produced for each scenario.

Figure 3. Land management allocation matrix.

Figure 4. The full 30 years of precipitation data plotted over time. First 20 years (red frame) are empirically measured rainfall events in Roujan (1992-2012), while the last 10 years are extension of the empirical dataset. The red star represents the amount of rainfall during the extreme event (435 mm/24h) measured in Perpignan in 1915.

Figure 5. Internal catchment dynamics under current land use and landscape structure (BLUS)
A: Land use and landscape structure, B-F: Erosion and deposition: (B) after 30 years of normal rainfall conditions, (C) after an extreme event in January, (D) after an extreme event in October. (E) internal catchment dynamics in year (indicated by number) when sediment export by the long-term rainfall series matches that of an extreme event in January, and (F) same as E, but for an extreme event in October. An elevation change legend (bottom right) shows deposition as positive, erosion as negative, and anything $<10^{-12}$m and $> -10^{-12}$m as “no erosion, no deposition”.


Figure 6. Internal catchment dynamics in one selected catchment segment
Land use and landscape structure(A), and erosion and deposition on different time scales (B-F) under different scenarios (indicated by rows) are shown in the selected catchment segment (locator figure, left bottom). Time scales: (B) 30 years under normal rainfall conditions, (C) an extreme event in January, and (D) an extreme event in October. In columns E and F, internal catchment dynamics are shown for the year when catchment outlet sediment export under long-term rainfall series met that of

an extreme event (E) in January and (F) in October.
Baseline scenario (B) represents steady production under current land use (LU) and landscape structure (S). More intensive (I) production than steady production is represented by ILU, ILS, ILUS scenario, less intensive (E) production by ELU, ELS, ELUS.
An elevation change legend (bottom right) shows deposition as positive, erosion as negative, and anything $<10^{-12}$m and $> -10^{-12}$m) as “no erosion, no deposition”.

Figure 7. External catchment dynamics

(A) Area specific sediment yield at the catchment outlet (10^{-2} Mg·km^{-2}) over 30 years. (B) Comparison between sediment delivery over long-term (30 years) and extreme events in January (arrow). Baseline scenario (B) represents steady production under current land use (LU) and landscape structure (S). More intensive (I) production than steady production is represented by ILU, ILS, ILUS scenario, less intensive € production by ELU, ELS, ELUS.

Figure 8. Land management effect on land degradation under different scenarios and temporal scales. Area-specific sediment budget within landscape management (10^{-2} Mg·km^{-2}) is plotted for long-term (30 years, x-axis), and an extreme event in January (y-axis). Erosion is plotted in red and orange, deposition in green and blue.





Baseline scenario: (A) land use and (B) landscape structure


- vineyards - mechanical weeding
- vineyards - chemical weeding
- annual crops - tillage
- fruit trees
- scrubland
- annual crops - no tillage
- pathway - bare soil
- pathway - vegetated soil
- grass strip
- building
- paved road

Management		Narrative		
		Stationary production (B)	More intensive production (I)	Less intensive production (E)
Landuse (LU)	Area: vineyards and crop	–	↑	↓
	Area: chemical weeding	–	↑	↓
	Area: no tillage	–	↓	↑
Landscape structure (LS)	Field unit surface	–	↑	↓
	Linear structure density	–	↓	↑
	Downslope tillage direction	–	↑	↑
	Area: grass strips	–	↓	↑


Scenarios were build according to three narratives: (B) – stationary production (representing current conditions), (I) more intensive production and (E) less intensive production than current conditions. Land use (LU) and/or landscape structure (LS) management was modified by increase (↑) or decrease (↓) in total area, and change in spatial allocation of LU and/or LD management in comparison with stationary production (-). The abbreviations of resulting seven scenarios (BLUS, ILU, ILS, ILUS, ELU, ELS, ELUS) indicate the narrative (B, I, S) and management (LU, LS, or LUS- both LU and LS). Detailed rules on allocation are in David et al. (2014). Spatial allocation of BLUS is plotted in Figure 5A (BLUS), and remaining scenarios in Supplementary Info 1.1A-1.6A.



Digital Elevation Model
Dynamic soil surface properties
Temporal dynamics:
Rainfall
Long-term (30 years)
Event occurring in
January
October
Land Management
Narrative
Stationary production (B)
More intensive production (I)
Less intensive production (E)
Modified
Land Use (LU)
Landscape structure (LS)
LU & LS
Land Use (LU)
Landscape structure (LS)
LU & LS
Scenarios:
BLUS
ILU
ILS
ILUS
ELU
ELS
ELUS
INPUTS
LANDSOIL
OUTPUTS
Spatial dynamics
Internal catchment dynamics (v)
(Spatial distribution of soil loss)
External catchment dynamics (∧)
(Sediment export at catchment outlet)



450
400
350
300
250
200
150
100
50
0
Rainfall events (mm)
0
20
30 years














Land use and landscape structure


- vineyards - mechanical weeding
- vineyards - chemical weeding
- annual crops - tillage
- fruit trees
- scrubland
- annual crops - no tillage
- pathway - bare soil
- patheway - vegetated soil
- grass strip
- building
- paved roads


Erosion

- < 0.3 m
- 0.01 to 0.3 m
- 0 to 0.01 m
- no erosion, no deposition

Accumulation

- 0 to 0.01 m
- 0.01 to 0.3 m
- > 0.3 m


A
B
C
D
E
F
BLUS
27 years
2 years
ILU
7 years
5 years
ILS
17 years
2 years
ILUS
6 years
4 years
ELU
24 years
3 years
ELS
26 years
2 years
ELUS
21 years
4 years
Selected area
Land use and landscape structure
vineyards - mechanical weeding
vineyards - chemical weeding
annual crops - tillage
fruit trees
scrubland
annual crops - no tillage
pathway - bare soil
pathway - vegetated soil
m
0 190 380
grass strip
building
paved road
Erosion
< 0.3 m
0.01 to 0.3 m
0 to 0.01 m
no erosion, no deposition
Accumulation
0 to 0.01 m
0.01 to 0.3 m
> 0.3 m


ILS
ILU
ILUS
ELS
ELU
ELUS
BLUS
A
Sediment export at catchment outlet (10-2 Mg·km-2)
120
100
80
60
40
20
0
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30
years
B
30
25
20
15
10
5
0
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 years
ILUS
ILU
ILS
BLUS
ELU
EQV
ELS
ELUS
Sediment export by extreme event in January (10-2 Mg·km-2)



BLUS
ILU
ILS
ILUS
ELU
ELS
ELUS
Extreme event (January)
Long-term
vineyard: weeding:
mechanical
chemical
crop:
tillage
no tillage
soil strip:
vegetated
unvegetated
scrubland
grass strip
fruit tree
road

Table 1. Mean soil loss for the catchment and soil export at catchment outlet under each scenario

	Long-term erosion			Extreme event in							
				January				October			
	Mean soil loss*	Soil export at outlet		Mean soil loss*	Soil export at outlet		Equal **	Mean soil loss*	Soil export at outlet		Equal **
	10^{-3}m	Mg	10^{-2} Mg·km^{-2}	10^{-3}m	Mg	10^{-2} Mg·km^{-2}	years	10^{-3}m	Mg	10^{-2} Mg·km^{-2}	years
BLUS	0.69±38.97	856.11	9.02	0.62±2.71	762.10	8.03	26.7	0.04±1.55	46.16	0.49	1.6
ILU	7.05±80.47	8700.59	91.70	1.62±4.39	1996.81	21.05	6.9	1.18±4.38	1456.43	15.35	5.0
ILS	1.73±35.69	2135.83	22.51	0.99±2.16	1216.83	12.82	17.1	0.09±0.74	112.82	1.19	1.6
ILUS	9.17±52.68	11309.58	119.20	1.89±3.44	2334.57	24.60	6.2	1.17±3.26	1445.42	15.23	3.8
ELU	0.62±25.16	769.80	8.11	0.50±2.01	614.55	6.48	23.9	0.07±1.16	88.06	0.93	3.4
ELS	0.38±35.01	469.59	4.95	0.33±3.10	412.83	4.35	26.4	0.03±1.38	36.87	0.39	2.4
ELUS	0.36±19.43	449.81	4.74	0.26±2.47	318.81	3.36	21.3	0.05±1.35	57.60	0.61	3.8

*- mean calculated based on raster cells, ± indicates standard deviation. Mean soil loss is also referred to as internal catchment dynamics in the text. Soil export is also referred to as external catchment dynamics in the text. **- number of years in which catchment outlet sediment export over long-term rainfall matched that of extreme events; BLUS-steady production - baseline land use and landscape structure. I-more intensive production than B. E-less intensive production than B; 10^{-2} Mg·km^{-2} equals t/ha