



HAL
open science

Could CTA be a reliable alternative method to identify non-genotoxic substances in Food Contact Material extract ?

Théo Lacour, Ruzanna Hayrapetyan, Isabelle Séverin, Isabelle Mouche, Francis Finot, Marie-Christine Chagnon

► To cite this version:

Théo Lacour, Ruzanna Hayrapetyan, Isabelle Séverin, Isabelle Mouche, Francis Finot, et al.. Could CTA be a reliable alternative method to identify non-genotoxic substances in Food Contact Material extract ?. ICT 2022 The XVIth International Congress of Toxicology, Sep 2022, Maastricht, Netherlands. , 368, pp.S141, 2022, 10.1016/j.toxlet.2022.07.402 . hal-03819008

HAL Id: hal-03819008

<https://hal-agrosup-dijon.archives-ouvertes.fr/hal-03819008>

Submitted on 18 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Could CTA be a reliable alternative method to identify non-genotoxic substances in Food Contact Material extract ?

T.Lacour^{1,2} and R. Hayrapetyan³, I. Severin³, I. Mouche¹, F. Finot¹, M.-C. Chagnon³

1 GenEvolutionN, Porcheville, France 2 Ecole Pratique des Hautes Etudes, Paris, France
3 Derttech "Packtox", NUTOX, INSERM U1231, Dijon, France


Introduction

The CTA using Bhas 42 cells, is a 3T3 mouse fibroblast cell line transfected with several copies of the Ras gene allowing a cell transformation process. Indeed the process of cellular transformation is possible because it is a cell already initiated.


This process is described by the formation of foci characterized by :

- deep basophilic staining
- spindle-shaped cells
- multilayer growth (piling up of cells)
- random cell orientation
- invasive cell growth into the background monolayer.

Transformed foci :



Non-transformed foci :



Non-intentionally added substances (NIAS) are chemical compounds that are present in food contact materials (FCMs) and could therefore migrate into food, but they are not added for a technical reason during the production process. Often their presence is not known by the consumer and not even by the producer.


Cell Transformation Assay (CTA) is able to detect non-genotoxic carcinogen agents but is it able to identify Non-Intentionally Added Substances (NIAS) ?

In this work the objective is to miniaturize the classical CTA. In a second step, the aim is to establish a more specific and sensitive labeling of transformed foci. Finally we would like to identify transcriptomics markers impacted by a non-genotoxic cancerigen agent.

Results

Labelling with BMVC

Transformed foci Non-transformed foci




Identification of the foci is done by an experienced experimenter. This staining allows an easier identification of the foci because it is realized under fluorescence.

In parallel, a quantitative examination can be performed. The fluorescent properties of BMVC allow a spectrophotometric determination with excitation at 435 nm and emission at 560 nm.

Labelling BMVC cells treated with caffeine or TPA

Promotion test

Initiation test




BMVC : 3,6-Bis(1-methyl-4-vinylpyridinium)-carbazole diiodide
TPA : 12-O-tetradecanoylphorbol-13-acetate

Unpaired t test with Welch's correction, **** : P value < 0,0001

Transcriptomics Assay

A transcriptomics assay allows the examination of the transcriptome to target which steps of the cell cycle are impacted by the test product.



Edited by Mascolo and al, 2018,

Previous publications illustrate its interest:

- HAPs (2, 4-diaminotoluene, benzo(a)pyrene, 2-acetylaminoflourene, or 3-methylcholanthrene) :
 - 14 genes regulated in common for the 24h and 32 days groups which shows the predictivity of this test
- 38 chemicals (tumor promoters, non-tumor promoters, genotoxic carcinogens and food additives) :
 - The cells are treated for 48 hours, resulting in only a list of genes commonly known to be involved in human cancers.

Conclusion

- The miniaturized 96-well CTA identifies genotoxic and non-genotoxic carcinogens using fewer cells, less medium and, most importantly, less test material.
- Labeling of BMVC on Bhas 42 cells combined with CTA would allow for more accurate and sensitive quantification, in addition to allowing for a non-examiner-dependent quantification.
- The transcriptome analysis allowed to identify the impacted cell cycle steps, such as immune response, cell adhesion, apoptosis and cell proliferation.

Outlook

- ❖ Automation of rinsing and identification of foci using a microplate washer and a microplate reader.
- ❖ Use BMVC labeling to identify the potential non-genotoxic carcinogenicity of NIAS.
- ❖ Reduction of SVF variability by using a serum substitute, non-fetal serum or by reducing the proportion of serum in the medium.
- ❖ Identification of non-genotoxic (epigenetic) damage with hypermethylation or hypomethylation as well as with different DNA compactions

References

- Maeshima H, Ohno K, Tanaka-Azuma Y, Nakano S, Yamada T. Identification of tumor promotion marker genes for predicting tumor promoting potential of chemicals in BALB/c 3T3 cells. *Toxicol In Vitro*. 2009 Feb;23(1):148-57. doi: 10.1016/j.tiv.2008.10.005. Epub 2008 Nov 1. PMID: 19000923.
- Mascolo MG, Perdichizzi S, Vaccari M, Rotondo F, Zanzi C, Grilli S, Paparella M, Jacobs MN, Colacci A. The transcriptomics assay: first steps for the development of an integrated approach to investigate the malignant cell transformation in vitro. *Carcinogenesis*. 2018 Jul 3;39(7):955-967. doi: 10.1093/carcin/bgy037. Erratum in: *Carcinogenesis*. 2018 Jul 3;39(7):968. PMID: 29554273; PMCID: PMC6031005.
- Rohrbeck A, Salinas G, Maaser K, Linge J, Salovaara S, Corvi R, Borlak J. Toxicogenomics applied to in vitro carcinogenicity testing with Balb/c 3T3 cells revealed a gene signature predictive of chemical carcinogens. *Toxicol Sci*. 2010 Nov;118(1):31-41. doi: 10.1093/toxsci/kfq246. Epub 2010 Aug 16. PMID: 20713471.